

University of Groningen

Immunologic aspects of ovarian cancer

Gooden, Marloes Jacomine Margaretha

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Gooden, M. J. M. (2013). *Immunologic aspects of ovarian cancer*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Chapter 3

Prognostic significance of tumor-infiltrating T-lymphocytes in primary and metastatic lesions of advanced stage ovarian cancer

Ninke Leffers*

Marloes J.M. Gooden*

Renske A. de Jong

Baukje-Nynke Hoogeboom

Klaske A. ten Hoor

Harry Hollema

H. Marike Boezen

Ate G.J. van der Zee

Toos Daemen

Hans W. Nijman

*Both authors contributed equally

Cancer Immunology Immunotherapy, 2009, 58: 449-459

Abstract

Purpose: Ovarian cancer patients with intra-tumoural CD3+ T-lymphocytes in primary tumour tissue have a better prognosis. This study aims to analyze the presence and relative influence of three important T-lymphocyte subsets, tumour-infiltrating CD8+ cytotoxic T lymphocytes (CTL), CD45R0+ memory T-lymphocytes and FoxP3+ regulatory T-lymphocytes (Treg), in primary tumour tissue and omental metastases of patients with ovarian cancer.

Experimental Design: The number of CD8+, CD45R0+ and FoxP3+ T-lymphocytes was determined by immunohistochemistry on a tissue micro array containing ovarian tumour tissue and/or omental metastases obtained at primary debulking surgery from 306 FIGO stage I-IV ovarian cancer patients. Immunohistochemistry data were correlated to clinicopathological parameters and survival data.

Results: High number of CD8+ CTL and a high CD8+/FoxP3+ ratio in ovarian-derived tumour tissue were associated with increased disease specific survival and proved to be independent prognostic factors in multivariate analyses. In advanced stage patients, the presence of CD8+ CTL, CD45R0+ memory T-lymphocytes, FoxP3+ Treg or a high CD8+/FoxP3+ ratio in ovarian-derived tumour tissue was associated with an increased disease-specific survival in univariate analysis, as was the presence of CD45R0+ memory T-lymphocytes and FoxP3+ Treg in omental metastases. Furthermore, in advanced stage patients CD8+ cytotoxic and FoxP3+ regulatory T-lymphocytes infiltrating ovarian-derived tumour tissue were independent predictors of increased prognosis.

Conclusions: T-lymphocytes infiltrating primary and metastatic ovarian cancer sites are associated with improved prognosis. These associations are especially distinct in advanced stage patients, underlining the potential for immunotherapy as a broadly applicable therapeutic strategy.

Introduction

Ovarian cancer is the most frequent cause of death from gynaecological malignancies, with an overall mortality of 60%. Due to its lack of specific symptoms, 70% of patients present with advanced disease. Initial treatment consists of cytoreductive surgery and platinum based chemotherapy. Even when initial treatment is successful, many patients relapse. There is no curative secondary treatment in case of residual disease after chemotherapy or relapse (1;2). Therefore, new therapeutic strategies are under investigation (3). Immunotherapy is one of these strategies and has yielded especially promising results in fundamental and animal research. However, clinical applications have only shown limited efficacy. Further knowledge is necessary to develop strategies to increase clinical efficacy of immunotherapy.

Tumour-infiltrating lymphocytes (TIL) were found to correlate with improved prognosis in several types of cancer, among which ovarian cancer (4-9). The presence of TIL is considered a reflection of the immune response to the tumour. However, not all T-lymphocyte subsets contribute equally to this observed positive effect on prognosis. Invasion by cytotoxic T lymphocytes (CTL) has been found to be advantageous in ovarian cancer (7) and several other types of cancer (10-13). The role of CD4+ T lymphocytes is ambiguous. Non-regulatory CD4+ helper T lymphocytes, especially of the Th1 subset, are considered to be beneficial in cancer in general (14). Regulatory CD4+ T-lymphocytes (Treg) have a physiological function in preventing autoimmunity. They may induce peripheral tolerance, and, in doing so, suppress immune responses. In cancer, Treg traffic to tumours as a result of chemokines produced by tumour cells and micro-environmental macrophages (15). There, they suppress effector T-lymphocytes by secreting transforming growth factor β (TGF- β) and interleukin 10 (IL-10) or by direct cell-cell contact (15). Treg have been found to unfavourably influence prognosis in ovarian cancer (15;16). From a functional point of view, the ratio between effector T-lymphocytes and regulatory T-lymphocytes may be even more interesting, as was demonstrated by Sato et al. (7) who showed that high ratios of CTL/helper T-lymphocytes and CTL/Treg were of prognostic significance, whereas no significant association between the number of tumour infiltrating T-lymphocytes in general or subsets (Treg / helper T-lymphocytes) and overall survival were found in ovarian cancer. A less well studied T-lymphocyte subset are memory T-lymphocytes, which may arise after the initial immune response against an antigen. In cancer, memory T-lymphocytes mediate long-term immunity against tumours (17). The presence of intra-tumoural memory T-lymphocytes is associated with increased survival in hepatocellular (18) and colorectal cancer (5;17). To our knowledge, no studies have been published investigating the presence of memory T-lymphocytes in ovarian cancer. In summary, patients with tumour-infiltrating T-lymphocytes mount an immune response against their tumour, the success of which seems to depend on the relative concentrations of different T-cell subtypes.

No reports have been published describing possible differences in T-lymphocyte infiltration patterns between primary tumour sites and metastases. The omentum is a frequent site for metastases in ovarian cancer, due to the displacement of tumour cells by the peritoneal fluid stream. We decided to study the presence of infiltrating T-lymphocytes in omental metastases, which was greatly

facilitated by the fact that omentectomy is a routine part of cytoreductive surgery. We compared the results with infiltrating T-lymphocytes in tissue, taken from the primary ovarian tumour.

Aims of the present study were to determine the presence of CD8+ CTL, CD45R0+ memory T-lymphocytes and FoxP3+ Treg in primary tumour specimens and simultaneous omental metastases of patients with ovarian cancer and to determine their prognostic impact.

Methods

Patients

As of 1985 the Department of Gynaecological Oncology at the University Medical Center Groningen (UMCG) keeps a computerized database of patients with malignant epithelial ovarian cancer treated at this hospital at any time point during the course of their disease, prospectively collecting information on clinicopathological characteristics and follow-up.

For this study, ovarian cancer patients were selected if primary surgery was performed by a gynaecological oncologist from the UMCG between May 1985 and June 2006 and if paraffin-embedded ovarian and omental tumour tissue was available.

Patients were staged according to FIGO classification (19). Tumours were graded and classified according to WHO criteria by a gynaecological pathologist (20). Adjuvant chemotherapy consisted of different platinum-based treatment regimens. Response to chemotherapy was evaluated according to WHO criteria (21). After treatment, patients were followed-up for at least 10 years with gradually increasing intervals. Follow-up data were completed for all patients until January 2007.

Institutional Review Board Approval

For the present study, all relevant data were retrieved from our computerized database into a separate anonymous database. In this separate, password-protected database, patient identity was protected by study-specific, unique patient codes, which were only known to two dedicated data managers, who also have daily responsibility for the larger database. In case of uncertainties with respect to clinicopathological and follow-up data, the larger databases could only be checked through the data managers, thereby ascertaining the protection of patients' identity. According to Dutch law no approval from our IRB was needed.

Tissue microarrays

Tissue microarrays (TMAs) were constructed as described in previous studies (22;23). In brief, paraffin-embedded tissue blocks containing tumour in ovarian and omental tissue, and corresponding haematoxylin & eosin (H&E) stained slides were retrieved from the pathology archives. Representative areas of tumour were marked on the H&E stained slides. Next, using these H&E slides for reference, four 0.6 mm core biopsies were taken from each tumour specimen and arrayed on a recipient paraffin block using a tissue microarrayer (Beecher instruments, Silver

Spring, Maryland, USA). Adhesion of cores to the recipient block was accomplished by placing the blocks in a 37°C oven for 15 min. For staining, 4µm sections were cut from each TMA block. H&E staining was performed to verify the presence of tumour in the arrayed samples.

Immunohistochemistry

TMA sections were stained with mouse monoclonal antibodies recognizing CTL (anti-CD8; Dako cytometry, Glostrup, Denmark), memory T-lymphocytes (anti-CD45R0 clone OPD4; Labvision, Fremont CA, USA) and Treg (anti-FoxP3 m22509, Abcam, Cambridge, UK). In brief, TMA sections were dewaxed in xylene and rehydrated using graded concentrations of ethanol to distilled water. After antigen retrieval, endogenous peroxidase activity was blocked by submersion of sections in a 0.3% H₂O₂ solution for 30 min. Sections were incubated with the primary antibody for 60 min at room temperature (dilutions: anti-CD8 1:20; anti-CD45R0 1:50; anti-FoxP3 1:100). Sections incubated with anti-CD45R0 and anti-FoxP3 were subsequently incubated with DAKO Envision+ for 30 min. For sections incubated with anti-CD8, RAMpo (dilution 1:100) and GARpo (dilution 1:100) were used as secondary and tertiary antibodies respectively. The antigen-antibody reactions were visualized with 3,3'-diaminobenzidine or NovaRED™ (Vector Laboratories, Burlingame) for anti-FoxP3 staining. Sections were counterstained with haematoxylin.




Figure 1 Immunohistochemical staining of tumor-infiltrating T-lymphocytes in ovarian cancer tissue. A. Flow diagram showing the type of tissue analyzed and the percentage of patients lost from analysis. Patients were excluded from analysis when less than two tissue cores with at least 20% tumor were present for evaluation. B. Intra-tumoral CD8+ T-lymphocytes, C. CD45R0+ T-lymphocytes, D. FoxP3+ T-lymphocytes at 400x magnification.

Table 1 Clinicopathological characteristics and survival data

	Total (n=306)		Primary ovarian cancer (n=270)		Omental metastases (n=147)	
<i>Age (years)</i>						
Mean (SD)	57.2	(13.5)	56.8	(13.8)	59.6	(12.2)
<i>DSS (months)</i>						
Median (95% C.I.)	37.4	(25.6-49.2)	45.4	(31.1-59.8)	20.4	(14.3-27.3)
<i>FIGO Stage</i>						
Stage I	67	(21.9%)	67	(24.8%)		
Stage II	24	(7.8%)	25	(9.3%)		
Stage III	171	(55.9%)	145	(53.7%)	116	(78.9%)
Stage IV	42	(13.7%)	32	(11.9%)	30	(20.4%)
Missing	2	(0.7%)	1	(0.4%)	1	(0.7%)
<i>Tumour type</i>						
Serous	171	(55.9%)	147	(54.4%)	105	(71.4%)
Mucinous	36	(11.8%)	35	(13.0%)	6	(4.1%)
Endometroid	41	(13.4%)	38	(14.1%)	13	(8.8%)
Clear Cell	21	(6.9%)	18	(6.7%)	7	(4.8%)
Adenocarcinoma	13	(4.2%)	10	(3.7%)	5	(3.4%)
Mixed Tumours	15	(4.9%)	14	(5.2%)	5	(3.4%)
Other	9	(2.9%)	8	(3.0%)	6	(4.1%)
<i>Tumour Grade</i>						
Grade I	52	(17.0%)	51	(18.9%)	6	(4.1%)
Grade II	80	(26.1%)	76	(28.1%)	28	(19.0%)
Grade III	135	(44.1%)	113	(41.9%)	91	(61.9%)
Undifferentiated	14	(4.6%)	11	(4.1%)	9	(6.1%)
Missing	25	(8.2%)	19	(7.0%)	13	(8.8%)
<i>Residual disease</i>						
< 2 cm	162	(52.9%)	157	(58.1%)	47	(32.0%)
>= 2 cm	123	(40.2%)	92	(34.1%)	94	(63.9%)
Missing	21	(6.9%)	21	(7.8%)	6	(4.1%)
<i>Chemotherapy</i>						
No chemotherapy	42	(13.7%)	40	(14.8%)	12	(8.2%)
Platinum-based	117	(38.2%)	104	(38.5%)	48	(32.7%)
Platinum & taxane based	113	(36.9%)	96	(35.6%)	69	(46.9%)
Other regimen	27	(8.8%)	24	(8.9%)	16	(10.9%)
Unknown	7	(2.3%)	6	(2.2%)	2	(1.4%)

DSS = disease-specific survival; FIGO = International Federation of Gynaecology and Obstetrics.

Scoring

All stainings were scored independently by two observers. Observers had no prior knowledge of clinicopathological information. For the anti-CD8 and anti-CD45RO staining, the number of cells with membrane staining within tumour epithelium was counted for each core. For the anti-FoxP3 staining, the number of cells with nuclear staining within tumour-islets in every core was counted.

To obtain a high concordance rate with whole tissue slides, we decided that minimally two cores with a minimum of 20% tumour tissue had to be present on the TMA for a sample to be entered into analysis (23). Subsequently, we calculated the number of intra-tumoural cells per 0,283 mm² of tumour (i.e. one whole core consisting of 100% tumour tissue), to correct for differences in the amount of tumour tissue and to standardize the analysis. This calculation entailed dividing the total number of intra-tumoural cells in cores containing $\geq 20\%$ of tumour by the total percentage of tumour tissue present in these cores.

Statistics

The distribution of all TIL subtypes was positively skewed. For further analysis, we therefore decided to categorize patients using the bottom tertile (p33) as a cut-off value (7). However, when the bottom tertile equalled zero, a subdivision based on the presence or absence of these cells was made for further analysis. Associations between clinicopathological characteristics and intra-tumoural T-lymphocytes were tested using the χ^2 test. Differences in numbers of tumour-infiltrating T-lymphocytes between primary ovarian tumour tissue and omental metastases were tested using Wilcoxon Signed Ranks test. Disease-specific survival (DSS) was defined as date of surgery until death due to ovarian cancer or date of last follow-up. DSS was estimated using Kaplan Meier and Log Rank test was used to assess for survival differences between groups. Cox proportional hazards models were used for multivariate analyses and were stratified for type of chemotherapy. Only variables that were significantly associated with DSS in the univariate analyses were entered into multivariate analyses. Survival analyses were performed in the total population as well as in a subgroup which included all advanced stage patients. For all tests, p-values < 0.05 were considered significant. All p-values were tested two-sided. All statistical analyses were performed using SPSS 14.0 software package for windows (SPSS Inc., Chicago, IL, USA).

Results

Study population

From a total of 306 ovarian cancer patients, sufficient paraffin-embedded ovarian and/or omental tissue containing epithelial ovarian carcinoma was available for construction of the TMA (figure 1a). Tumour-containing ovarian tissue was available from 270 patients. From 111 (41.1%) of these patients tumour-containing omental tissue was also available. Only tumour-containing omental tissue was available from an additional 36 patients. Clinicopathological characteristics of patients are summarized in Table 1. Forty-two patients did not receive chemotherapy, 28 of whom were diagnosed with FIGO stage I disease. The remaining 14 patients were either not fit or unwilling to receive chemotherapy. Of the patients treated with chemotherapy 89.5% received a platinum-based regimen. The median survival for all patients was 37.4 months with an estimated 5-year disease specific survival of 41%.

Table 2 Relationship of tumor-infiltrating lymphocytes in ovarian tissue to clinicopathological parameters and each other.

	CD8+ T lymphocyte			CD45RO+ T lymphocyte			FoxP3+ T lymphocyte		
	Lowest tertile (%)	All others (%)	p-value ^a	Absent (%)	Present (%)	p-value ^a	Absent (%)	Present (%)	p-value ^a
<i>Age (years)</i>									
<59	43 (38.7)	68 (61.3)	0.101	61 (55.0)	50 (45.0)	0.571	48 (44.9)	59 (55.1)	0.575
≥59	33 (28.4)	83 (71.6)		62 (51.2)	59 (48.8)		54 (48.6)	57 (51.4)	
<i>FIGO stage</i>									
Stage I/II	25 (33.3)	50 (66.7)	0.947	45 (59.2)	31 (40.8)	0.173	39 (57.4)	29 (42.6)	0.031
Stage III/IV	51 (33.8)	100 (66.2)		77 (49.7)	78 (50.3)		62 (41.6)	87 (58.4)	
<i>Tumor type</i>									
Serous	37 (30.3)	85 (69.7)	0.278	68 (52.7)	61 (47.3)	0.917	51 (42.1)	70 (57.9)	0.125
Non-serous	39 (37.1)	66 (62.9)		55 (53.4)	48 (46.6)		51 (52.6)	46 (47.4)	
<i>Differentiation grade</i>									
Grade I/II	39 (37.5)	65 (62.5)	0.208	56 (54.9)	46 (45.1)	0.513	55 (57.3)	41 (42.7)	0.011
Grade III & undiff.	32 (29.4)	77 (70.6)		57 (50.4)	56 (49.6)		43 (39.4)	66 (60.6)	
<i>Residual disease</i>									
<2cm	45 (36.0)	80 (64.0)	0.692	64 (48.9)	67 (51.1)	0.109	58 (50.4)	57 (49.6)	0.577
≥2cm	28 (33.3)	56 (66.7)		51 (60.0)	34 (40.0)		39 (46.4)	45 (53.6)	
<i>CD8+ T lymphocyte</i>									
Lowest tertile									
All others						<0.001	53 (75.7)	17 (24.3)	<0.001
<i>CD45RO+ T lymphocyte</i>									
Absent							47 (32.9)	96 (67.1)	
Present							66 (70.2)	41 (37.6)	<0.001
<i>CD8+/FoxP3+ ratio</i>									
Bottom tertile						0.038	28 (29.8)	68 (62.4)	
All others				41 (63.1)	24 (26.9)				
<i>CD8+/CD45RO+ ratio</i>				64 (47.4)	71 (52.6)				
Bottom tertile									
All others							50 (64.1)	28 (35.9)	<0.001
							43 (35.5)	78 (64.5)	

FIGO = International Federation of Gynaecology and Obstetrics. ^a P-values were calculated using Pearson Chi square-test. Bold signifies p<0.05.

Tumour-infiltrating T-lymphocytes

Intra-tumoural CD8+, CD45R0+ and FoxP3+ T-cells were present in 91.2%, 47.0% and 53.2% of primary ovarian tissue, respectively (representative examples in figure 1b-d), while intra-tumoural CD8+, CD45R0+ and FoxP3+ T-cells were present in 96.2%, 49.3% and 73.1% of omental metastases, respectively.

In both ovarian and omental derived tumour tissue, the number of tumour-infiltrating T-lymphocytes was positively associated with each other for all subtypes (Tables 2 and 3). Although the median number of tumour-infiltrating T-lymphocytes was higher for omental than ovarian tissue, no differences in intra-tumoural T-lymphocytes were observed between ovarian and omental tumour tissue within patients for whom both samples were available (data not shown).

Association of tumour-infiltrating lymphocytes with clinicopathological parameters

Table 2 and 3 show the relationship of clinicopathological parameters to the presence of intra-tumoural TIL in ovary and omentum derived tumour tissue respectively. In ovarian tissue, the presence of FoxP3+ cells was positively associated with advanced stage disease ($p=0.031$). Furthermore, patients with poorly differentiated tumours were more likely to have intra-tumoural FoxP3+ cells compared to patients with well-differentiated tumours ($p=0.011$). Also, patients younger than 59 years at time of diagnosis were more likely to have a low CD8+/CD45R0+ ratio than older patients (data not shown: 48.5% vs. 34.3%, $p=0.036$).

In omental metastases, high CD8+ T-cell numbers were associated with <2 cm residual disease after primary debulking surgery ($p=0.035$). A similar association was found for FoxP3+ cells, which were more often present in patients with residual disease of <2cm ($p=0.028$).

Survival analysis

In agreement with expectations, well-known prognostic factors such as age ≥ 59 years, advanced stage disease, poorly differentiated tumours, serous tumours and ≥ 2 cm residual disease after primary debulking surgery were associated with a shorter disease specific survival (data not shown). Furthermore, univariate analysis of disease specific survival showed an initial survival advantage for patients with high numbers of intra-tumoural CD8+ cytotoxic T-lymphocytes in ovarian-derived tumour tissue, which subsides after 10 years of follow-up ($p=0.042$, figure 2a). Disease-specific survival was not influenced by the presence of CD45R0+ memory T-lymphocytes, FoxP3+ regulatory T-lymphocytes in ovarian tissue. However, median DSS of patients with a high CD8+/FoxP3+ ratio in ovarian-derived tumour tissue was twice as high as for patients with a low CD8+/FoxP3+ ratio (50.0 vs. 23.0 months, $p=0.014$, figure 2b). For the CD8+/CD45R0+ ratio a trend was observed towards a longer median DSS in patients with a high ratio (51.7 vs. 30.9 months, $p=0.056$).

We subsequently repeated univariate survival analyses using a subgroup consisting of patients with advanced stage disease. Interestingly, in this subgroup a prolonged median DSS was observed for all T-lymphocyte subsets studied in ovarian-derived tumour tissue, except for the CD8+/CD45R0+ ratio (30.5 vs. 15.6 months, $p=0.076$), i.e. CD45R0+ lymphocytes ($n=155$, 37.3 vs. 16.4 months, $p=0.009$, figure 2c), CD8+ lymphocytes ($n=150$, 30.3 vs. 12.2 months, $p=0.012$, figure 2d), FoxP3+

Table 3 Relationship of intra-tumoral TIL in omental tissue to clinicopathological parameters and each other

	CD8+ T lymphocyte			CD45RO+ T lymphocyte			FoxP3+ T lymphocyte		
	Lowest tertile (%)	All others (%)	p-value ^a	Absent (%)	Present (%)	p-value ^a	Absent (%)	Present (%)	p-value ^a
Age (years)									
<59	28 (38.4)	45 (61.6)	0.173	38 (53.5)	33 (46.5)	0.495	28 (39.4)	43 (60.6)	0.653
≥59	16 (27.1)	43 (72.9)		30 (47.6)	33 (52.4)		21 (35.6)	38 (64.4)	
Tumor type									
Serous	32 (33.3)	64 (66.7)	1.000	52 (54.7)	43 (45.3)	0.149	38 (40.4)	56 (59.6)	0.299
Non-serous	12 (33.3)	24 (66.7)		16 (41.0)	23 (59.0)		11 (30.6)	25 (69.4)	
Differentiation grade									
Grade I/II	12 (41.4)	17 (58.6)	0.244	15 (51.7)	14 (48.3)	0.871	12 (42.9)	16 (57.1)	0.356
Grade III & undiff.	28 (29.8)	66 (70.2)		47 (50.0)	47 (50.0)		31 (33.3)	62 (66.7)	
Residual disease									
<2cm	7 (18.9)	30 (81.1)	0.035	18 (46.2)	21 (53.8)	0.488	8 (22.2)	28 (77.8)	0.028
≥2cm	34 (38.2)	55 (61.8)		47 (52.8)	42 (47.2)		38 (43.2)	50 (56.8)	
CD8+ T lymphocyte									
Lowest tertile									
All others						<0.001	29 (59.2)	14 (17.5)	<0.001
CD45RO+ T lymphocyte									
Absent									
Present							36 (73.5)	27 (34.6)	<0.001
CD8+/FoxP3+ ratio									
Bottom tertile						0.007	13 (26.5)	51 (65.4)	
All others				28 (66.7)	14 (33.3)				
CD8+/CD45RO+ ratio									
Bottom tertile				35 (41.2)	50 (58.8)				
All others							34 (59.6)	23 (40.4)	<0.001
							22 (27.5)	58 (72.5)	

FIGO = International Federation of Gynaecology and Obstetrics. ^a P-values were calculated using Pearson Chi square-test. Bold signifies p<0.05.

lymphocytes (n=149, 30.3 vs. 14.0 months, $p=0.008$, figure 2e) and a high CD8+/FoxP3+ ratio (n=146, 30.3 vs. 14.6 months; $p=0.016$, figure 2f). Similarly, univariate analysis of disease-specific survival differences based on the presence of TIL in omental metastases, which are by definition present only in patients with advanced stage disease, showed an increased survival in patients with intra-tumoural FoxP3+ Treg (25.6 vs. 14.7 months, $p<0.008$, figure 2g) and CD45RO+ memory T-lymphocytes (24.3 vs. 15.6 months, $p=0.031$, figure 2h).

Multivariate analysis was performed stratified for type of chemotherapy. For all patients as well as the advanced stage subgroup, the above-mentioned well-known prognostic factors and tumour-infiltrating T-lymphocytes were entered into the Cox proportional hazards model (table 4). In the analysis using all patients, CD8+ lymphocytes and CD8+/FoxP3+ ratio were independent prognostic factors for DSS (HR 0.36, 95% C.I. 0.22-0.59, $p<0.001$; resp. HR 0.53, 95% C.I. 0.32-0.85, $p=0.009$), next to commonly acknowledged risk factors as stage of disease, differentiation grade and amount of residual tumour after primary debulking surgery. When analyzing only patients with advanced stage disease, a high number of CD8+ cytotoxic T-lymphocytes in ovarian-derived tumour tissue was demonstrated to be an independent prognostic factor for longer DSS (HR 0.35, 95% C.I. 0.21-0.60, $p<0.001$), as was the presence of FoxP3+ Treg (H.R. 0.55, 95% C.I. 0.34-0.88, $p=0.013$). Additionally, multivariate analysis was performed using the absolute number of tumour-infiltrating T-lymphocytes instead of the dichotomized variables (data not shown). Unlike CD8+ lymphocytes, FoxP3+ lymphocytes infiltrating tumour epithelium remained an independent factor for longer DSS (HR 0.937, 95% C.I. 0.88-1.00, $p=0.035$).

Discussion

The observation of an improved survival of patients with intra-tumoural T-lymphocytes suggests an important role for the immune system in the natural course of ovarian cancer (9). In the present study, we analyzed the relative contribution of different tumour-infiltrating T-lymphocyte subtypes to this observed improvement of survival. To our knowledge, this is the first study to examine the prognostic influence of different tumour-infiltrating T-lymphocytes in primary as well as metastatic lesions. We show improved disease-specific survival in advanced stage ovarian cancer patients with CD45RO+ memory T-lymphocytes and/or FoxP3+ regulatory T-lymphocytes infiltrating ovarian-derived tumour tissue and/or omental metastases. In addition, we also confirm important previous observations by others of improved survival in patients with either a high number of tumour-infiltrating CD8+ cytotoxic T-lymphocytes or a high CD8+/FoxP3+ ratio (7).

Intra-tumoural CD45RO+ T-lymphocytes, present in approximately 50% of patients, were associated with increased DSS when present in ovarian-derived tumour tissue and omental metastases. Although its expression is not restricted to memory T-lymphocytes, CD45RO is commonly used as a marker for memory T-lymphocytes (5;17;24). Upon antigenic stimulation, naïve T-lymphocytes differentiate into effector T-lymphocytes, but also memory T-lymphocytes can be formed. The latter assure a strong and speedy immune response on subsequent exposures to the same antigen.




Figure 2 Disease-specific survival (in months) of ovarian cancer patients based on tumor-infiltrating lymphocytes. Cumulative survival time was estimated by the Kaplan–Meier method. Log Rank test was used to evaluate survival differences between groups. High numbers of CD8+ T-lymphocytes (a) and a high CD8+/Foxp3+ ratio (b) in ovarian derived tumor tissue were associated with improved survival in FIGO stage I–IV disease. In advanced stage patients, a survival benefit was observed for patients with CD45RO+ T-lymphocytes (c) CD8+ T-lymphocytes (d), FoxP3+ T-lymphocytes (e) a high CD8+/Foxp3+ ratio (f) in ovarian-derived tumor tissue. Furthermore, FoxP3+ (g) and CD45RO+ T-lymphocytes (h) in omental metastases were associated with improved disease-specific survival.

The presence of tissue-infiltrating memory T-lymphocytes thus implies prior exposure to an antigen encountered in the tissue environment, which in case of tumour-infiltrating memory T-lymphocytes may very well reflect an anti-tumour response of the immune system. The presence of a survival benefit for patients with high levels of infiltrating memory T-lymphocytes was previously described in colorectal cancer (5;17). The cell numbers seen in these studies were remarkably higher than those used in our study (cut-off values 80-250 cells/mm²), which may be explained in several ways. Firstly, no distinction was made between tumour- and peri-tumoural stroma infiltrating cells in these studies, whereas we only report tumour-infiltrating memory T-lymphocytes. Secondly, when

Table 4 Multivariate Cox regression analysis on disease-specific survival.

All stages	H.R.	95% CI	p-value	H.R.	95% CI	p-value
Age ≥ 59 years	0.96	0.60-1.55	0.876	0.98	0.61-1.59	0.938
FIGO Stage II/IV	3.31	1.34-8.13	0.009	3.59	1.73-9.37	0.009
Grade II/undifferentiated	1.97	1.19-3.27	0.008	1.64	0.98-2.74	0.059
Non-serous tumour	0.67	0.41-1.12	0.125	0.63	0.38-1.05	0.078
Residual tumour	3.38	2.27-6.44	<0.001	3.01	1.82-4.96	<0.001
CD8+ T-lymphocytes high (OV)	0.36	0.22-0.59	<0.001			
CD8+/FoxP3+ ratio high (OV)				0.53	0.32-0.85	0.009
Advanced stage only	H.R.	95% CI	p-value	H.R.	95% CI	p-value
Age ≥ 59 years	0.82	0.50-1.36	0.444	1.02	0.63-1.64	0.945
Grade II/undifferentiated	1.75	1.02-2.99	0.041	1.82	1.07-3.09	0.026
Non-serous tumour	0.76	0.44-1.30	0.309	0.77	0.44-1.33	0.346
Residual tumour	3.58	2.08-6.17	<0.001	2.41	1.50-3.88	<0.001
CD8+ T-lymphocytes high (OV)	0.35	0.21-0.60	<0.001			
CD45RO+ T-lymphocytes present (OV)				0.72	0.46-1.14	0.166
FoxP3+ T-lymphocytes present (OV)				0.55	0.34-0.88	0.013
Advanced stage only, cont.	H.R.	95% CI	p-value	H.R.	95% CI	p-value
Age ≥ 59 years	1.00	0.61-1.65	1.000	0.76	0.43-1.33	0.888
Grade II/undifferentiated	1.42	0.83-2.44	0.205	1.04	0.57-1.89	0.910
Non-serous tumour	0.76	0.44-1.30	0.318	0.60	0.34-1.06	0.078
Residual tumour	2.50	1.51-4.15	<0.001	2.63	1.41-4.90	0.002
CD8+/FoxP3+ ratio high (OV)	0.68	0.42-1.08	0.099			
CD45RO+ T-lymphocytes present (OM)				0.69	0.43-1.08	0.106
FoxP3+ T-lymphocytes high (OM)						

Analyses were performed stratified for type of chemotherapy. FIGO = International Federation of Gynecology and Obstetrics, H.R. = hazard ratio; C.I. = confidence interval; OV = ovarian-derived tumour tissue; OM = omental metastasis; Bold signifies p<0.05.

repeating our analysis on the total cell count/core we still did not observe levels similar to that in colorectal cancer, although results for the survival analysis were comparable to the analysis with tumour-infiltrating memory T-lymphocytes only (data not shown). A more likely explanation is that the higher number of infiltrating lymphocytes in colon tissue could be considered inherent to its daily exposure to foreign products and pathogens, an observation which does not hold for ovarian tissue.

Our data show that patients with intra-tumoural CTL infiltrations and/or a high CTL/Treg ratio in tumour-containing ovarian tissue consistently have the longest disease-specific survival, confirming the results previously described in a smaller population of ovarian cancer patients (7). However, unlike previous studies in ovarian cancer addressing the prognostic influence of Treg (7;15), we found tumour-infiltrating Treg to be an independent positive factor for disease-specific survival in patients with advanced stage disease. One possible explanation is the fact that although FoxP3 is currently the best marker available for immunohistochemical staining of Treg on paraffin embedded tissue, it is not exclusively expressed in regulatory cells. Recently, it was shown that activated CD4+CD25- effector T cells transiently express FoxP3, without having regulatory activity (25;26). Possibly, the positive prognostic effect of the presence of FoxP3+ cells in our samples can be attributed to the staining of not only suppressive, but also activated T-lymphocytes expressing FoxP3. Similar observations of positive effects of Treg have however previously been reported in studies on immune cell malignancies and head and neck squamous cell carcinoma (HNSCC), some of which further phenotyped the FoxP3+ cells. A high number of Treg is associated to longer survival in follicular lymphoma and classic Hodgkin's lymphoma (27;28). In HNSCC where higher numbers of circulating Treg were present in patients without evidence of disease after primary treatment compared to patients with active disease, loco-regional Treg were associated with better loco-regional control (29;30). Furthermore, the possible protective role for Treg against cancer was also illustrated in murine models of colorectal cancer, which show that adoptive transfer of CD4+CD25+ regulatory cells results in regression of adenomas (31;32). Although we find that the presence of Treg is associated with an increase in disease-specific survival, we also find that 1) the absolute number of Treg present does not seem to alter this positive prognostic effect, 2) the presence of Treg is strongly associated with high numbers of CTL, and 3) a high CTL/Treg ratio is associated with a longer disease-specific survival. It thus seems plausible that it is not merely the presence of Treg, but rather the infiltration of tumour by T-lymphocytes in general which is responsible for the observed survival advantage, as was previously reported by Zhang et al. (9). In this point of view, infiltration of tumour epithelium, or any other inflammatory site, by lymphocytes is naturally accompanied by regulatory T-cells and the effectiveness of immune responses depends on the proportion of the different lymphocyte subtypes present instead of on the presence of a particular subtype.

Unlike previous studies of TIL in ovarian cancer, we have also studied the prognostic influence of TIL in omental metastases of ovarian cancer patients. Although TIL in omental metastases were not independent predictors of disease-specific survival, we did show an increased survival in univariate analysis for advanced stage patients with memory T-lymphocytes and Treg infiltrating

omental metastases. Interestingly, median numbers of TIL were higher in omental metastases than in ovarian-derived tumour tissue and the percentage of patients with Treg in omental metastases was substantially higher than for ovarian-derived tumour tissue. The omentum may be considered a secondary lymphoid tissue as it contains large aggregates of macrophages and lymphocytes, the so-called milky spots. These milky spots contain mostly macrophages (68%), but T-lymphocytes are also present (10%) (33). The high percentage of patients with Treg observed in omental metastases and the higher median cell counts as compared to ovarian-derived tumour tissue may be a reflection of the lymphoid function of the omentum.

In summary, we show that memory T-lymphocytes infiltrating ovarian-derived tumour tissue and omental metastases are associated with increased disease specific survival in patients with advanced stage disease. Unexpectedly, the presence of regulatory T-lymphocytes in ovarian-derived tumour tissue is also an independent factor for increased disease specific survival. However, the fact that a high CTL/Treg ratio independently predicts increased survival as well, suggests that it is not so much the presence of Treg as the presence of TIL in general that is responsible for the observed survival effect. Our results support the notion that different T-lymphocyte subsets play a unique and important role in the natural course of ovarian cancer.

References

1. Greenlee RT, Hill-Harmon MB, Murray T, Thun M. Cancer statistics, 2001. *CA Cancer J Clin* 2001 Jan;51(1):15-36.
2. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2005 Mar;55(2):74-108.
3. Agarwal R, Linch M, Kaye SB. Novel therapeutic agents in ovarian cancer. *Eur J Surg Oncol* 2006 May 13;32(8):875-86.
4. Clemente CG, Mihm MC, Jr., Bufalino R, Zurrida S, Collini P, Cascinelli N. Prognostic value of tumor infiltrating lymphocytes in the vertical growth phase of primary cutaneous melanoma. *Cancer* 1996 Apr 1;77(7):1303-10.
5. Galon J, Costes A, Sanchez-Cabo F, Kirilovsky A, Mlecnik B, Lagorce-Pages C, et al. Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* 2006 Sep 29;313(5795):1960-4.
6. Halpern AC, Schuchter LM. Prognostic models in melanoma. *Semin Oncol* 1997 Feb;24(1 Suppl 4):S2-S7.
7. Sato E, Olson SH, Ahn J, Bundy B, Nishikawa H, Qian F, et al. Intraepithelial CD8+ tumor-infiltrating lymphocytes and a high CD8+/regulatory T cell ratio are associated with favorable prognosis in ovarian cancer. *Proc Natl Acad Sci U S A* 2005 Dec 20;102(51):18538-43.
8. Vesalainen S, Lipponen P, Talja M, Syrjanen K. Histological grade, perineural infiltration, tumour-infiltrating lymphocytes and apoptosis as determinants of long-term prognosis in prostatic

adenocarcinoma. *Eur J Cancer* 1994;30A(12):1797-803.

9. Zhang L, Conejo-Garcia JR, Katsaros D, Gimotty PA, Massobrio M, Regnani G, et al. Intratumoral T cells, recurrence, and survival in epithelial ovarian cancer. *N Engl J Med* 2003 Jan 16;348(3):203-13.
10. Kondratiev S, Sabo E, Yakirevich E, Lavie O, Resnick MB. Intratumoral CD8+ T lymphocytes as a prognostic factor of survival in endometrial carcinoma. *Clin Cancer Res* 2004 Jul 1;10(13):4450-6.
11. Naito Y, Saito K, Shiiba K, Ohuchi A, Saigenji K, Nagura H, et al. CD8+ T cells infiltrated within cancer cell nests as a prognostic factor in human colorectal cancer. *Cancer Res* 1998 Aug 15;58(16):3491-4.
12. Nakano O, Sato M, Naito Y, Suzuki K, Orikasa S, Aizawa M, et al. Proliferative activity of intratumoral CD8(+) T-lymphocytes as a prognostic factor in human renal cell carcinoma: clinicopathologic demonstration of antitumor immunity. *Cancer Res* 2001 Jul 1;61(13):5132-6.
13. Schumacher K, Haensch W, Roefzaad C, Schlag PM. Prognostic significance of activated CD8(+) T cell infiltrations within esophageal carcinomas. *Cancer Res* 2001 May 15;61(10):3932-6.
14. Yu P, Fu YX. Tumor-infiltrating T lymphocytes: friends or foes? *Lab Invest* 2006 Mar;86(3):231-45.
15. Curiel TJ, Coukos G, Zou L, Alvarez X, Cheng P, Mottram P, et al. Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival. *Nat Med* 2004 Sep;10(9):942-9.
16. Wolf D, Wolf AM, Rumpold H, Fiegl H, Zeimet AG, Muller-Holzner E, et al. The expression of the regulatory T cell-specific forkhead box transcription factor FoxP3 is associated with poor prognosis in ovarian cancer. *Clin Cancer Res* 2005 Dec 1;11(23):8326-31.
17. Pages F, Berger A, Camus M, Sanchez-Cabo F, Costes A, Molitor R, et al. Effector memory T cells, early metastasis, and survival in colorectal cancer. *N Engl J Med* 2005 Dec 22;353(25):2654-66.
18. Cai XY, Gao Q, Qiu SJ, Ye SL, Wu ZQ, Fan J, et al. Dendritic cell infiltration and prognosis of human hepatocellular carcinoma. *J Cancer Res Clin Oncol* 2006 May;132(5):293-301.
19. Cancer Committee of the International Federation of Gynaecology and Obstetrics. Staging announcement: FIGO Cancer Committee. *Gynecol Oncol* 1986;25:383-5.
20. Scully RE. *Histological Typing of Ovarian Tumours*. 2 ed. Berlin: Springer; 1999.
21. World Health Organization. *Handbook for Reporting Results of Cancer Treatment*. Geneva: World Health Organization; 1979.
22. Kononen J, Bubendorf L, Kallioniemi A, Barlund M, Schraml P, Leighton S, et al. Tissue microarrays for high-throughput molecular profiling of tumor specimens. *Nat Med* 1998 Jul;4(7):844-7.
23. Rosen DG, Huang X, Deavers MT, Malpica A, Silva EG, Liu J. Validation of tissue microarray technology in ovarian carcinoma. *Mod Pathol* 2004 Jul;17(7):790-7.
24. Anraku M, Cunningham K, Yun Z, Tsao M, Zhang L, Keshavjee S, et al. Impact of tumor-infiltrating T cells on survival in patients with malignant pleural mesothelioma. *The Journal of Thoracic and Cardiovascular Surgery* 2008 Apr;135(4):823-9.
25. Allan S, Crome S, Crellin N, Passerini L, Steiner T, Bacchetta R, et al. Activation-induced FOXP3

in human T effector cells does not suppress proliferation or cytokine production. *Int Immunol* 2007 Apr 1;19(4):345-54.

26. Roncarolo MG, Gregori S. Is FOXP3 a bona fide marker for human regulatory T cells? *Eur J Immunol* 2008 Apr;38(4):925-7.
27. Alvaro T, Lejeune M, Salvado MT, Bosch R, Garcia JF, Jaen J, et al. Outcome in Hodgkin's lymphoma can be predicted from the presence of accompanying cytotoxic and regulatory T cells. *Clin Cancer Res* 2005 Feb 15;11(4):1467-73.
28. Lee A, Clear A, Calaminici M, Davies A, Jordan S, MacDougall F, et al. Number of CD4+ Cells and Location of Forkhead Box Protein P3-Positive Cells in Diagnostic Follicular Lymphoma Tissue Microarrays Correlates With Outcome. *J Clin Oncol* 2006 Nov 1;24(31):5052-9.
29. Badoual C, Hans S, Rodriguez J, Peyrard S, Klein C, Agueznay NH, et al. Prognostic value of tumor-infiltrating CD4+ T-cell subpopulations in head and neck cancers. *Clin Cancer Res* 2006 Jan 15;12(2):465-72.
30. Strauss L, Bergmann C, Gooding W, Johnson J, Whiteside T. The Frequency and Suppressor Function of CD4+CD25highFoxp3+ T Cells in the Circulation of Patients with Squamous Cell Carcinoma of the Head and Neck. *Clin Cancer Res* 2007 Nov 1;13(21):6301-11.
31. Erdman S, Poutahidis T, Tomczak M, Rogers A, Cormier K, Plank B, et al. CD4+ CD25+ Regulatory T Lymphocytes Inhibit Microbially Induced Colon Cancer in Rag2-Deficient Mice. *Am J Pathol* 2003 Feb 1;162(2):691-702.
32. Erdman S, Sohn J, Rao V, Nambiar P, Ge Z, Fox J, et al. CD4+CD25+ Regulatory Lymphocytes Induce Regression of Intestinal Tumors in ApcMin/+ Mice. *Cancer Res* 2005 May 15;65(10):3998-4004.
33. Krist LF, Eestermans IL, Steenbergen JJ, Hoefsmit EC, Cuesta MA, Meyer S, et al. Cellular composition of milky spots in the human greater omentum: an immunochemical and ultrastructural study. *Anat Rec* 1995 Feb;241(2):163-74.

